

A clinical case of feline crown restoration with monolithic zirconia

Sinem COŞKUN^{1,a,✉}, Çağlar COŞKUN^{2,b}

¹Lokman Hekim University, Faculty of Dentistry, Oral and Maxillofacial Radiology Department, Ankara, Türkiye; ²Veterinary Practice Dost Pati, Ankara, Türkiye

^aORCID: <https://orcid.org/0000-0003-4772-6047>; ^bORCID: <https://orcid.org/0000-0002-3265-969X>

ARTICLE INFO

Article History

Received : 24.08.2023

Accepted : 01.06.2024

DOI: 10.33988/auvfd.1349278

Keywords

Animal dentistry

Root canal treatment

Veterinary dentistry

✉Corresponding author

sinem.coskun@lokmanhekim.edu.tr

How to cite this article: Coşkun S, Coşkun Ç (XXXX): A clinical case of feline crown restoration with monolithic zirconia. Ankara Univ Vet Fak Derg, XX (X), 000-000. DOI: 10.33988/auvfd.1349278.

ABSTRACT

Direct pulp exposure necessitates immediate extraction or root canal treatment in pets. Monolithic zirconia is used as a crown material in dentistry. In the current study, a 4-year-old female Ankara cat had a monolithic zirconia crown restored following root canal therapy. The tooth underwent root canal therapy, and preparation was done for a fixed monolithic zirconia prosthetic full crown restoration. Monolithic zirconia crowns are new, significant restorative materials that have been used rarely and little studied in veterinary dentistry despite having important material properties. Consequently, this is the first reported case of monolithic zirconia crown restoration from Türkiye, and the use of this material is strongly suggested to clinicians.

Direct pulp exposure from a fractured tooth occurs in 10% of pet animals, and it necessitates immediate extraction or root canal treatment (RCT) (1). Patients who have been exposed to dentine because of trauma from falling, being hit by cars, or fighting should have a radiograph taken to check for signs of infection, inflammation, or loss of tooth vitality (3, 9). Bacterial invasion through the dentinal tubules may result in pulpitis. The only appropriate clinical solution is immediate RCT and restoration in one visit (13). Monoblock yttrium oxide stabilized tetragonal zirconia polycrystals are widely used by clinicians in fixed crown restorations due to their biomechanical advantages, such as high mechanical properties, dimensional and chemical stability, fracture resistance, and non-toxic properties (4,5). Monolithic zirconia (MZ) ceramic is the restoration of only zirconia ceramic in one piece (6). Since it does not contain a superstructure, it has high fracture resistance (8). The aim of this study is to present the endodontic treatment of pulp exposed after trauma in pet animals and its restoration with MZ crowns.

A complicated left maxillary canine tooth fracture due to falling on a 4-year-old female cat was diagnosed. An oral and dental examination confirmed the tooth as suffering from a significant crown fracture with pulp exposure under sedation (Figure 1a). Enamel tissue loss necessitated a full crown restoration and RCT. A standard RCT was performed with complete isolation using a rubber dam under general anesthesia. The tooth was ultrasonically scaled prior to the RCT. Handheld endodontic files and rotary devices were used to clean, disinfect, and shape the root canal before sealing the pulp cavity. Sodium hypochlorite and EDTA solutions were used as lubricants and endodontic irrigants to disinfect the canal. Several files were used to remove the exposed pulp through the coronal fracture site (Figure 1b). Before and immediately after the treatment, digital radiographic images of the affected tooth were taken using the bisecting angle technique in order to determine the working length and check the final obturation (Figure 1c). The root canals were dressed after being dried with paper points. A lentulo

spiral filler was used to introduce endodontic sealer. At the 12,5 mm working length, calibrated gutta-percha points were positioned and compressed vertically without heating the compacting tool. The canal was filled consistently. Self-adhesive resin cement was used to lute the glass fiber post. The core was constructed from composite resin. The tooth was prepared following polymerization (Figure 1d). With the putty wash technique, impressions were made. A three-dimensional computer-aided design technology was used to produce the plaster model. The restoration was made from a MZ semi-sintered ceramic block using computer-aided manufacturing. Examining and making the necessary adjustments to the internal adaptation of the crown on the model, occlusal contacts, marginal fit, harmony of the external contours, and approximate surfaces. Afterwards, external B1 color staining and glaze were done. In the patient's mouth, the restoration was evaluated. The crown was cemented with self-adhesive resin cement. The restoration was exposed to the curing light for 20 seconds. Overflowing cement residues were cleaned immediately (Figure 1e). Approximately one month after the procedure, the patient returned to the referring veterinarian for a postoperative check-up. The gingiva in the area was in good health and showed no signs of

inflammation. The MZ crown fit perfectly. The patient had resumed regular chewing patterns and was eating normally. About six months after the initial treatment, the patient went back to the referring vet for a conscious oral examination. The tooth and its surrounding structure healed properly. Palpation revealed no signs of pain, and the tooth remained functional. There was no swelling, movement, or fluctuation during the intraoral inspection. The marginal adaptation of crown restoration was well done. On probing, there were no signs of caries. Moreover, the restoration still has its original hue. No gingival recession occurred. The teeth of the antagonist have no wear (Figure 1f).

Canine teeth commonly encounter dentin and enamel breaking off since they are the longest and most front teeth (10). RCT is frequently used on pets to avoid tooth loss (11, 12). In this case, a complicated crown fracture was determined, and a single-session RCT was applied. Prosthodontic crowns have been recorded in client-owned animals during the past 45 years as a technique to support endodontic therapy, replace missing tooth structure, safeguard the tooth from future harm, and finally restore the patient's normal function (12, 13). Despite the fact that full metal crowns are commonly used in pets, there is limited research in the literature using MZ crowns (9, 14).




Figure 1. a: Pulp exposure of left maxillary canine tooth b: The initial file for the tooth. c: The final obturation. d: Tooth preparation for monolithic zirconia crown. e: Monolithic zirconia crown. f: Clinical appearance of six months follow-up.

Coffman et al. (5) mentioned zirconia crowns as an alternative to porcelain or ceramic restorations. Ahmed et al. (2) applied two different crowns and found that apoptosis occurred less with zirconia crowns. Similar to this case, Fink et al. (7) reported two cases of tooth fractures restored with zirconia crowns after RCT to protect teeth from further damage. After being informed of their options for care, the owners chose to proceed with a RCT and subsequent prosthetic repair. The findings of this case study lead us to the conclusion that MZ crown restorations following RCT demonstrated adequate strength and could serve as a replacement for complete metal crown restorations in animal dentistry. Further studies are needed to assess long-term efficacy and treatment success in other animal species.

Acknowledgments

We would like to thank dentist Orhan Coşkun for his valuable contributions to the study.

Financial Support

This research received no grant from any funding agency/sector.

Conflict of Interest

The authors declared that there is no conflict of interest.

Author Contributions

Both SC and ÇC conceived and planned the study. SC and ÇC did clinical and radiographic examination, patient's follow-up. All authors contributed to the interpretation of the results. SC took the lead in writing the manuscript. All authors provided critical feedback and helped shape the research, analysis and manuscript.

Data Availability Statement

The data supporting this study's findings are available from the corresponding author upon reasonable request.

Ethical Statement

This clinical case report does not present any ethical concerns.

Animal Welfare

The authors confirm that they have adhered to ARRIVE Guidelines to protect animals used for scientific purposes.

References

1. **Abdulmajeed AA, Lim KG, Närhi TO, et al** (2016): *Complete-arch implant-supported monolithic zirconia fixed dental prostheses: A systematic review*. J Prosthet Dent, **115**, 672-677.
2. **Ahmed RK, Korsel AM, Elshahawy W** (2018): *Gingival Response To Zirconia And Porcelain Fused To Metal Crowns: An Experimental Study*. Egypt Dent J, **64**, 1973-1981.
3. **Bellows J** (2004): The Comprehensive Oral Prevention, Assessment and Treatment Visit. 221-237. In: IA Ames(Ed), Small Animal Dental Equipment, Materials, and Techniques: A Primer. Wiley Blackwell Pub.
4. **Charnock M, Usher FJ** (1968): *Fitting a prosthetic crown to a canine tooth in a police dog*. Vet Rec, **83**, 464-465.
5. **Coffman C, Visser C, Soukup J, et al** (2019): Crowns and prosthodontics. 387-410. In: HB Lobprise, RD Jonathan (ed), Wiggs's Veterinary Dentistry: Principles and Practice. 2nd ed. John Wiley & Sons, Pondicherry, India.
6. **Çelik M, Bural C, Bayrakdar G** (2014): *Diş Hekimliğinde Zirkonya Uygulamaları*. Atatürk Üniversitesi Diş Hekimliği Fakültesi Dergisi, **24**, 106-116.
7. **Fink L, Reiter AM** (2015): *Assessment of 68 prosthodontic crowns in 41 pet and working dogs (2000–2012)*. J Vet Dent, **32**, 148-154.
8. **Guess PC, Bonfante EA, Silva NR, et al** (2013): *Effect of core design and veneering technique on damage and reliability of Y-TZP-supported crowns*. Dent Mater, **29**, 307-316.
9. **Mele RE, Caiafa A, Kurtzman GM** (2017): *Feline dental implants: long-term follow-up of two cases*. J Vet Dent, **34**, 268-278.
10. **Nakamura K, Harada A, Kanno T, et al** (2015): *The influence of low-temperature degradation and cyclic loading on the fracture resistance of monolithic zirconia molar crowns*. J Mech Behav Biomed Mater, **47**, 49-56.
11. **Ribka EP, Niemiec BA** (2021): *Success of Feather Margin Preparation for Full Metal Prosthodontic Crowns in the Canine Teeth in 84 Pet and Working Dogs (2005-2017)*. J Vet Dent, **39**, 34-40.
12. **Soukup JW, Hetzel S, Paul A** (2015): *Classification and epidemiology of traumatic dentoalveolar injuries in dogs and cats: 959 injuries in 660 patient visits (2004–2012)*. J Vet Dent, **32**, 6-14.
13. **Strøm PC, Arzi B, Lommer MJ, et al** (2018): *Radiographic outcome of root canal treatment of canine teeth in cats: 32 cases (1998–2016)*. J Am Vet Med Assoc, **252**, 572-580.
14. **Wingo K** (2018): *Cementation of full coverage metal crowns in dogs*. J Vet Dent, **35**, 46-53.

Publisher's Note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.